

# New records of predation and feeding behavior of the vine snake *Chironius brazili* (Squamata: Colubridae) in the Espinhaço Mountain Range, Brazil

Vitor Otávio Ferreira Assunção, Gabriel Saturno

<sup>1</sup> Research Laboratory in Herpetology, Institute of Biosciences, Federal University of Mato Grosso do Sul, Campo Grande, Mato Grosso do Sul 79070-900, Brazil

Recibida: 21 Marzo 2025  
Revisada: 09 Junio 2025  
Aceptada: 10 Julio 2025  
Editor Asociado: V. Arzamendia

doi: 10.31017/CdH.2025.(2025-010)

## ABSTRACT

The vine snake *Chironius brazili* inhabits rocky *campo rupestre* environments in the Espinhaço Mountain Range, Brazil. Although species of *Chironius* primarily feed on anurans, we report new prey records for *C. brazili*, including the Cycloramphidae and the first documented case of fish predation in the genus. These records expand the known diet of *C. brazili* and demonstrate its ability to exploit different environments while foraging.

Key Words: : Snake, Diet, Cycloramphidae, Fish.

The vine snake *Chironius brazili* Hamdan and Fernandes, 2015, is found in the ecoregions of the Cerrado, Atlantic Forest, and Pampa, primarily inhabiting gallery forests and open areas (Hamdan and Fernandes, 2015; Ugalde *et al.*, 2024). Species of the genus *Chironius* Fitzinger, 1826, exhibit morphology primarily adapted for an arboreal lifestyle (*i.e.* slender body, long tail, and large number of vertebrae; Banci *et al.*, 2022; Parreira *et al.*, 2024) and have a predominantly batrachophagous diet, although they also feed on reptiles, mammals, and birds (Roberto and Souza, 2020). However, *C. brazili* can exploit different substrates while foraging, including rocky outcrops along waterfall margins in mountainous regions of the Brazilian Cerrado (Parreira *et al.*, 2024).

The Espinhaço Mountain Range, in eastern Brazil, extends for approximately 1200 km across the states of Minas Gerais and Bahia, with altitudes ranging from 800 to 2000 m a.s.l. (Silveira *et al.*, 2016).

This mountain range forms a transition between the Atlantic Forest to the east and the open landscapes of the Cerrado and Caatinga to the north and west (Guedes *et al.*, 2020; Santana *et al.*, 2023). Above 1000 m a.s.l., *campo rupestre* (rupestrian fields) predominates, an environment characterised by shrub-grass vegetation interspersed with rocky outcrops and open areas with sandy soil and low water availability (Silveira *et al.*, 2016; Santana *et al.*, 2023).

Although the low tree cover density in the open areas of the Cerrado and *campo rupestre*, *Chironius* species occurring in these environments primarily feed on hylids (Kok, 2010; Fernandes and Hamdan, 2014; Hamdan and Fernandes, 2015; Roberto and Souza, 2020; Parreira *et al.*, 2024). This association suggests that, even in predominantly open habitats, *Chironius* maintains its dietary preference for arboreal anurans (Roberto and Souza, 2020; Parreira *et al.*, 2024). This study presents new observations


on the feeding and hunting behavior of *C. brazili* in the *campo rupestre*, highlighting a new predation record on the Cycloramphidae big-eared river frog, *Thoropa megatympanum* Caramaschi and Sazima, 1984, and the first documented case of fish predation in *Chironius*.

On 16 October 2022, at approximately 03:40 pm, in the southern region of the Espinhaço Mountain Range, in the municipality of Presidente Kubitschek, Minas Gerais, Brazil ( $18^{\circ}27'16.7''\text{S}$   $43^{\circ}36'10.9''\text{W}$ ) (Fig. 1), an adult *C. brazili* was observed foraging on a rocky outcrop, moving through crevices and cavities on the rock surface, close water bodies and waterfalls, as reported by Parreira *et al.* (2024). The snake inserted its head into a rocky foramen, struck, and captured an adult *T. megatympanum* (Fig. 2), extracting it from the rock crevices and swallowing it. After digestion, the snake resumed its foraging activity, climbed onto the rocks, and captured another *T. megatympanum*, repeating the same capture and consumption process. Both anurans

were ingested in a postero-anterior direction within approximately 20 minutes.

The second record was made on 10 October 2024, at approximately 02:20 pm, also in the southern region of the Espinhaço Mountain Range, in the municipality of Diamantina, Minas Gerais, Brazil ( $18^{\circ}11'01.08''\text{S}$ ,  $43^{\circ}37'05.60''\text{W}$ ) (Fig. 1). An adult *C. brazili* was observed foraging within a waterfall, near rocky outcrops. During the observation, the snake submerged in the water and, while still underwater, it grabbed an unidentified fish, which it carried to the surface and swam towards one of the banks surrounding the waterfall. This is the first known record of *Chironius* preying on a fish.

Information on the foraging and diet of *C. brazili* is scarce (Roberto and Souza, 2020; Parreira *et al.*, 2024). Until now, only two prey species have been recorded for this snake, both occurring in high-altitude rocky streams being *Bokermannohyla martinsi* (Bokermann, 1964) ( $n=1$ ) and *B. pseudopseudis* (Miranda-Ribeiro, 1937) ( $n=3$ ) (Lucas *et al.*, 2017;



**Figure 1.** (A) Topographic map showing the known distribution of vine snake *Chironius brazili* (black circles), big-eared river frog *Thoropa megatympanum* (white circles) and our records (blue triangles). (B) Enlarged region showing the location of predation events in the southern portion of the Espinhaço Mountain Range (Minas Gerais, Brazil).





**Figure 2.** Adulte vine snake *Chironius brazili* with an adult big-eared river frog *Thoropa megatympanum* in its mouth in Espinhaço Mountain Range (Minas Gerais, Brazil).

Parreira *et al.*, 2024). Similarly, *T. megatympanum* also inhabits high-altitude *campo rupestre* environments (Fig. 1B), using these habitats for shelter and reproductive vocalization (Caramaschi and Sazima, 1983; Sabbag *et al.*, 2018). The association of these anurans with streams, rocky outcrops, and sparse vegetation makes them potential prey for snakes that forage in such environments (e.g., Roberto and Souza, 2020; Parreira *et al.*, 2024). Although the diet of the genus *Chironius* is largely based on hylid frogs (Roberto and Souza, 2020), our records indicate that *C. brazili* exhibits an opportunistic feeding strategy, consuming the prey available in its habitat. This report expands the knowledge of the species diet and highlights its efficient foraging capability in rocky Cerrado environments. Moreover, the predation of a fish by this species suggests a greater dietary flexibility, as well as the ability to explore aquatic environments in search of prey.

## Acknowledgments

The authors would like to thank the Ericles Silva and João Vítor Pinheiro dos Santos for having recorded the events. This study was financially supported by the Federal University of Mato Grosso do Sul – UFMS/MEC. Assunção, V. and Saturno, G. thanks the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brazil (CAPES) – Funding code 001.

## Literature cited

- Banci, K.R.D.S.; Guimarães, M.; Siqueira, L.H.C.; Muscat, E.; Sazima, I. & Marques, O.A.V. 2022. Body shape and diet reflect arboreality degree of five congeneric snakes sympatric in the Atlantic forest. *Biotropica* 54: 839-851.
- Caramaschi, U. & Sazima, I. 1983. Uma nova espécie de *Thoropa* da Serra do Cipó, Minas Gerais, Brasil (Amphibia, Leptodactylidae). *Revista Brasileira de Zoologia* 2: 139-146.
- Fernandes, D.S. & Hamdan, B. 2014. A new species of *Chironius* Fitzinger, 1826 from the state of Bahia, Northeastern Brazil (Serpentes: Colubridae). *Zootaxa* 3881: 563-575.
- Guedes, T.B.; Azevedo, J.A.R.; Bacon, C.D.; Provete, D.B. & Antonelli, A. 2020. Diversity, endemism, and evolutionary history of montane biotas outside the Andean region. In *Neotropical diversification: Patterns and processes* (pp. 299-328).
- Hamdan, B. & Fernandes, D.S. 2015. Taxonomic revision of *Chironius flavolineatus* (Jan, 1863) with description of a new species (Serpentes: Colubridae). *Zootaxa* 4012 (1): 97-119.
- Kok, P.J.R. 2010. A new species of *Chironius* Fitzinger, 1826 (Squamata: Colubridae) from the Pantepui region, northeastern South America. *Zootaxa* 2611: 31-44.
- Lucas, D.F.; Silva, D.; Rocha, P.; Barbosa, A. & Feio, R.N. 2017. Predação do hílideo *Bokermannohyla martinsi* (Bokermann, 1964) pela serpente *Chironius brazili* (Hamdan & Fernandes, 2015) na Reserva Particular do Patrimônio Natural Santuário do Caraça, Catas Altas, Minas Gerais, Brasil. In *VIII Congresso Brasileiro de Herpetologia*, Campo Grande, Brasil.
- Parreira, I.; Kimura, P.C.; Carvalho Lima, L.F.; de Freitas, M.A. & Brandão, R.A. 2024. A new foraging habitat for *Chironius* vine snakes and a new prey for *Chironius brazili* in open rocky Cerrado habitats in Central Brazil. *Studies on Neotropical Fauna and Environment* 59: 659-663.
- Roberto, I.J. & Souza, A.R. 2020. Review of prey items recorded for snakes of the genus *Chironius* (Squamata, Colubridae), including the first record of *Osteocephalus* as prey. *Herpetology Notes* 13: 1-5.
- Sabbag, A.F.; Lyra, M.L.; Zamudio, K.R.; Haddad, C.F.B.; Feio, R.N.; Leite, F.S.F.; Gasparini, J.L. & Brasileiro, C.A. 2018. Molecular phylogeny of Neotropical rock frogs reveals a long history of vicariant diversification in the Atlantic forest. *Molecular Phylogenetics and Evolution* 122: 142-156.
- Santana, D.J.; Ragalzi, E.; Koroiva, R.; Mângia, S.; Ceron, K.; Leite, F.S. & Shepard, D.B. 2023. Lineage diversification of the Sky Island treefrog *Scinax curicica* (Anura, Hylidae) in the Espinhaço Mountain range. *Biological Journal of the Linnean Society* 142: 58-67.

V. Assunção & G. Saturno — Predation behavior of *Chironius brazili*

Silveira, F.A.O.; Negreiros, D.; Barbosa, N.P.U.; Buisson, E.; Carmo, F.F.; Carstensen, D.W.; Conceição, A.A.; Cornelissen, T.G.; Echternacht, L.; Fernandes, G.W.; Garcia, Q.S.; Guerra, T.J.; Jacobi, C.M.; Lemos-Filho, J.P.; Le Stradic, S.; Morellato, L.P.C.; Neves, F.S.; Oliveira, R.S.; Schaefer, C.E.; Viana, P.L. & Lambers, H. 2016. Ecology and evolution of plant diversity in the endangered *campo*

*rupestre*: A neglected conservation priority. *Plant and Soil* 403: 129-152.

Ugalde, M.R.; Guimarães, M.R.S.; Luna, I.V.; Guedes, T.B. & Hamdan, B. 2024. First record of *Chironius brazili* Hamdan & Fernandes, 2015 (Squamata, Colubridae) for the state of Tocantins and north region of Brazil. *Herpetologia Brasileira* 13: 140-146.

© 2025 por los autores, licencia otorgada a la Asociación Herpetológica Argentina. Este artículo es de acceso abierto y distribuido bajo los términos y condiciones de una licencia Atribución-No Comercial 4.0 Internacional de Creative Commons. Para ver una copia de esta licencia, visite <http://creativecommons.org/licenses/by-nc/4.0/>